## What is claimed is:

SUBA,>

5

- An antimicrobial copolymer obtainable by copolymerizing (component I) aliphatically unsaturated monomers which have been functionalized by means of an ester group and at least singly functionalized by means of a tertiary amino group with (component II) another aliphatically unsaturated monomer which has been at least singly functionalized by means of an amino group, where component I and component II are different from one another.
- 10 2. The antimicrobial copolymer as claimed in claim 1, wherein

component II is composed of aliphatically unsaturated monomers which have been at least singly functionalized by means of a tertiary amino group.

15 3. The antimicrobial copolymer as daimed in claim 1 or 2, wherein

component I is composed of aliphatically unsaturated monomers whose ester group has been at least singly functionalized by means of an amino group.

- 4. The antimicrobial copolymer as claimed in one of claims 1 to 3, wherein component I is composed of acrylate or methacrylates which have been at least singly functionalized by means of a tertiary amino group.
- 25 5. The antimicrobial polymer as claimed in one of claims 1 to 4,
  wherein
  each of components I and II is an aliphatically unsaturated monomer
  functionalized by means of a tertiary amino group and having the general formula

 $R^{1}NR^{2}R^{3}$ 

where R<sup>1</sup>: is a branched, unbranched or cyclic, saturated or

SSESHYS NIESD

30

unsaturated hydrocarbon radical having up to 50 carbon atoms which may have substitution by O atoms, N atoms or S atoms, and

R<sup>2</sup> and R

5

are branched, unbranched or cyclic, saturated or unsaturated hydrocarbon radicals having up to 25 carbon atoms, which may have substitution by O atoms, N atoms or S atoms, where R<sup>2</sup> and R<sup>3</sup> are identical or different,

with the proviso that R<sup>1</sup> in monomers of component I contains an ester group.

The antimicrobial coating made from antimicrobial copolymers as claimed in one 10 6. of claims 1 to 5,

wherein

the copolymerization is carried oùt on a substrate.

The antimicrobial coating made from antimicrobial copolymers as claimed in one 15 7. of claims 1 to 5,

wherein

the copolymerization is carried out as a graft polymerization of a substrate.

8. The antimicrobial coating as claimed in claim 7, wherein

20

Ú

the substrate is activated prior to the graft polymerization by UV radiation, plasma treatment, corona treatment, flame treatment, ozonization, electrical discharge or γ-radiation.

25

9. The antimicrobial coating as claimed in claim 7, wherein the substrate is activated prior to the graft polymerization by UV radiation with a photoinitiator.

A process for preparing antimicrobial copolymers by copolymerizing (component I) aliphatically unsaturated monomers which have been functionalized by means

of an ester group and a tertiary amino group with (component II) another aliphatically unsaturated monomer which has been at least singly functionalized by means of an amino group, where components I and II are different from one another.

5

11. The process as claimed in claim 10.

wherein

component it is composed of aliphatically unsaturated monomers which have been at least singly functionalized by means of a tertiary amino group.

10

The process as claimed in claim 10 or 11, 12.

wherein

component I is composed of aliphatically unsaturated monomers whose ester group has been at least singly functionalized by means of an amino group.

DSGEB476 GIRSIE 15

The process as claimed in one of claims 10 to 12,

wherein

component I is composed of acrylate or methacrylates which have been at least singly functionalized by means of a tertiary amino group.

20

14. The process as claimed in one of claims 10 to 13,

wherein

each of components I and II is an allehatically unsaturated monomer functionalized by means of a tertiary amino group and having the general formula

25

## R<sup>1</sup>NR<sup>2</sup>R<sup>3</sup>

where R1: is a branched, unbranched or cyclic, saturated or unsaturated hydrocarbon radical having up to 50 calbon atoms which may have substitution by O atoms, N atoms or S atoms, and

R<sup>2</sup> and R<sup>3</sup>:

are branched, unbranched or cyclic, saturated or unsaturated hydrocarbon radicals having up to 25 carbon atoms, which

30

15

20

25

30

may have substitution by O atoms, N atoms or S atoms, where R<sup>2</sup> and R<sup>3</sup> are identical or different,

with the proviso that R<sup>1</sup> in monomers of component I contains an ester group.

5 15. The process as chaimed in one of claims 10 to 14, wherein

the copolymerization is carried out on a substrate.

- 16. The process as claimed in one of claims 10 to 15,
- 10 wherein

the copolymerization is carried out as a graft polymerization of a substrate.

17. The process as claimed in claim 16,

wherein

the substrate is activated prior to the graft polymerization by UV radiation, plasma treatment, Corona treatment, flame treatment, ozonization, electrical discharge or  $\gamma$ -radiation.

ENOH2

18. The process as claimed in claim 17,

wherein

the substrate is activated prior to the graft polymerization by UV radiation with a photoinitiator.

- 19. The use of the antimicrobial copolymers as claimed in one of claims 1 to 9 for producing products with an antimicrobial coating comprising the copolymer.
  - 20. The use of the antimicrobial polymers as claimed in one of claims 1 to 9 for producing items in medical technology with an antimicrobial coating comprising the copolymer.
  - 21. The use of the antimicrobial copolymers as claimed in one of claims 1 to 9 for producing hygiene items with an antimicrobial coating comprising the copolymer.

22. The use of the antimicrobial copolymers as claimed in one of claims 1 to 9 in surface coatings, protective paints or in other coatings.

APPH3